

## Hungarian contribution to the mineralogy and geology of clays

Commemorating the 50<sup>th</sup> anniversary of the Hungarian Clay Minerals Group founded in 1960

István VICZIÁN

Geological Institute of Hungary, 1143 Budapest, Stefánia út 14., Hungary (viczian@mafi.hu)

### *Magyar eredmények az agyagásványtan és -földtan terén*

*Az 1960-ban alapított MFT Agyagásványtani Szakosztály 50. évfordulója alkalmából*

#### Összefoglalás

A cikk a magyar földtudományi agyagkutatás főbb eredményeit foglalja össze témák szerint csoportosítva.

**Általános művek.** A Magyar honi Földtani Társulat Agyagásványtani Szakosztályának alapítója, NEMECZ írta az első magyar általános szakkönyvet az agyagásványokról. Tágabb értelemben BÁRDOSSY alapvető műveit a bauxitokról szintén az agyagkutatás körébe sorolhatjuk. A talajok agyagásványaival STEFANOVITS, SZENDREI és NEMECZ, újabban pedig NAGY és KÓNYA könyvei foglalkoztak.

**Egyes agyagásványok vizsgálata.** A Füzérradványon bányászott híres illites ásványt MAEGDEFRAU és HOFMANN írta le még 1937-ben. Ezt nevezhetjük „a világ második illitjének” (az illinoisi után). Az ásvány további kutatásáról NEMECZ és VARJÚ számolt be. A vastartalmú agyagásványok, mint a szeladonit, glaukonit és a „Fe-dús montmorillonit csoport” kémiai rendszerezését lényegesen előrevitték WEISZBURG és TÓTH E. munkái. ERDÉLYI részletesen tanulmányozott számos szerpentinásványt. Újabban DÓDONY végzett szerkezeti vizsgálatokat szerpentinásványokon (pl. antigorit, poligonális szerpentin).

**Agyagásványok meghatározási módszerei.** Új, és ma is használható röntgendiffrakciós mennyiségi határozási módszert NÁRAY-SZABÓ vezetett be. A PAULIK testvérvármű termikus meghatározó készüléket fejlesztett ki, amelyet „derivatográf”-nak neveztek. Az első termikus határozó atlaszt az agyagásványokra vonatkozólag FÖLDVÁRI-VOGL állította össze. A termikus módszert jelenleg is magas szinten alkalmazza FÖLDVÁRI.

**Talajok és üledékek.** Magyarország talaj-agyagásvány térképét STEFANOVITS és DOMBÓVÁRI készítették el. BIDLÓ részletesen vizsgálta a vörös agyagokat, amelyek legtöbbször fosszilis talajokból származnak. Bazaltkráterekben fosszilis bentonitos tavi üledékeket ismert fel SOLTI. Ezek anyagát FÖLDVÁRI, Zs. BARNA és mások vizsgálták. A Kárpát-medence főleg riolitos eredetű bentonittelepeit KOVÁCS-PÁLFFY vizsgálja rendszeresen.

**Közetrétegtani alkalmazások.** A magyarországi üledékes formációkat agyagásvány-tartalmuk szempontjából VICZIÁN jellemzette. Jelenleg ezt a kutatási irányt RAUCSIK, VARGA A. és SZAKMÁNY folytatják tovább. VICZIÁN, MÁTYÁS J. és mások a szmektit–illit diagenetikus átalakulást alkalmazták a szénhidrogén-prognózisban. Az igen kisfokú metamorfózis rétegszilikájnainak fő kutatója ÁRKAI, munkatársai TÓTH M., JUDIK és mások. Ő vezette be a klorit „kríztalysossági” fokot (Árkai-index), munkatársaival vizsgálta az indexásványokat, a kőzetek szövétét stb. Kutatásaikat Magyarország határain túlra, a svájci Alpokra és DK-Európára is kiterjesztették.

A hidrotermális elváltozási öveket a Tokaji-hegységben elsőnek SZÉKYNÉ FUX ismerte fel. Később NEMECZ és VARJÚ, újabban MOLNÁR részletesen vizsgálták az elváltozással létrejött agyagásványokat. A nemesagyag-telepek kutatásával évtizedekig MÁTYÁS E. és ZELENKA foglalkozott, alkalmazhatóságukat és kolloid tulajdonságaiat JUHÁSZ, SZÁNTÓ és tanítványaik vizsgálták. A nehézfémek által okozott környezeti szennyezés modellezését NÉMETH végzi. SZABÓ egyik fő szakértője az agyagos szigetelőréteg kialakításának városi hulladéklerakók körül. Agyagos mellékkőzetben kívánják elhelyezni a nagy intenzitású radioaktív hulladékot a Mecsek hegységben.

Tárgyszavak: agyagásványok, szedimentológia, talajtan, igen kisfokú metamorfózis, környezeti ásványtan

#### Abstract

The main results of the Hungarian clay research are reviewed in the field of earth sciences in a thematic order.

A general textbook on clay minerals was published by the founder of the Clay Minerals Group, Professor NEMECZ. In a broader sense, the important books on bauxites by BÁRDOSSY belong to the field of clay research. Soil clay minerals were discussed in the textbooks of STEFANOVITS, SZENDREI and NEMECZ, and recently by NAGY and KÓNYA.

**Specific clay minerals.** The famous illite mineral of Füzérradvány — the “second illite in the world” (after the illite of

Illinois) — was described by MAEGDEFRAU & HOFMANN as early as 1937 and later reviewed by NEMECZ & VARJÚ. WEISZBURG and E. TÓTH contributed much to the chemical systematics of celadonite, glauconite and the “Fe-rich montmorillonite” series. Earlier, serpentine minerals were extensively studied by ERDÉLYI and in recent years by DÓDONY (structural analyses of antigorite and polygonal serpentine).

*Determinative methods in clay mineralogy.* A method of quantitative analysis by X-ray diffraction was introduced by NÁRAY-SZABÓ. A special apparatus for thermal analysis — called the “Derivatograph” — was developed by PAULIK et al. An atlas of thermal analysis was first compiled by FÖLDVÁRI-VOGL. The application of this method was later continued at a higher level by FÖLDVÁRI.

*Soils and sediments.* STEFANOVITS and DOMBÓVÁRI published a map of clay minerals in Hungarian soils. Red clays, mostly relict fossil soils were studied by BIDLÓ. FEKETE published a book on recent tropical soils. Fossil bentonitic lake sediments of basaltic craters were recognised by SOLTI and analysed by FÖLDVÁRI, Zs. BARNA and others. KOVÁCS-PÁLFFY focused his systematic studies on rhyolite-related bentonite deposits of the Carpathian Basin.

*Lithostratigraphic applications.* Clay minerals of Hungarian sedimentary formations were systematically characterised by VICZIÁN. Today this line has been continued by RAUCSIK, A. VARGA and SZAKMÁNY. The diagenetic transformation of smectite to illite was widely applied for CH prospecting by VICZIÁN, J. MÁTYÁS and others. Layer silicates of the very lowgrade metamorphic stage have been studied by ÁRKAI and his co-workers M. TÓTH, JUDÍK etc. He introduced the parameter “chlorite crystallinity” (Árkai index), studied index minerals, and also applied textural analysis. His studies extended to Hungary, the Swiss Alps, and to South East Europe.

Zones of hydrothermal alteration in the Tokaj Mts were first recognised by SZÉKY-FUX and later intensively studied by NEMECZ and VARJÚ and recently by MOLNÁR. Clay deposits of economic value were studied for several decades by E. MÁTYÁS and ZELENKA, their application and colloidal properties were analysed by JUHÁSZ, SZÁNTÓ and their students. Model experiments of environmental pollution caused by heavy metals were carried out by NÉMETH. Clay liners for communal waste depositories were developed by SZABÓ. Pelitic host rocks for high intensity radioactive waste depositories have been studied in Mecsek Mts.

**Keywords:** clay mineralogy, sedimentology, soil science, very low-grade metamorphism, environmental mineralogy

## Introduction

The Hungarian Clay Minerals Group was founded in 1960. In this review, on the occasion of the 50th anniversary of this Group, the main results of Hungarian clay research in the field of earth sciences are presented. With respect to the references in given here, only a few publications have been selected. The main considerations were that the work is typical for the author or the subject. This historical account follows the example of the review made by VOGL (1982) on the development of the analytical methods during the first 20 years of the Clay Group (1960–1980). A valuable source of data regarding the activity in the early periods can be found in the bibliography compiled by SZENDREI (1979). This bibliography is available in manuscript form and contains nearly 500 items. A similar bibliographic compilation, with brief historical notes, was written by VICZIÁN in 1999. Recently valuable bibliographical and biographical data were collected by SZENDREI (2010) for the purposes of a home page of the Hungarian Geological Society.

## Books on Clay Minerals

A general textbook on clay minerals was published by the founder of the Clay Mineral Group, Professor NEMECZ (1981). In a broader sense the important books of BÁRDOSSY (1982, BÁRDOSSY & ALEVA 1990) on bauxites and the bauxite sedimentological studies of MINDSZENTY (1999) belong to the clay research. Soil clay minerals were discussed in the textbooks of STEFANOVITS (1981), SZENDREI (1994) and NEMECZ (2006). FEKETE (1988) published a book on tropical soils. It is not intended here to review the colloid chemistry of

clay minerals in detail. Only a few university courses and textbooks on colloids which are closely related to clay minerals are mentioned. Such are the university courses by JUHÁSZ (1995–1996), the book by SZÁNTÓ (1987) and the recently published work by M. NAGY & KÓNYA (2009). JUHÁSZ & OPOCZYK (1982, 1990) have become international-renowned specialists on the mechanochemistry of silicate minerals.

## Specific clay minerals

The famous illite mineral of the locality of Füzérradvány, — also called sarospatakite and more recently “Zempleni illite” — can be regarded as the “second illite of the world” because it was described by MAEGDEFRAU & HOFMANN as early as 1937 (after the Illinois illite: see VICZIÁN 2000). Later the mineral was reviewed by NEMECZ & VARJÚ (1970) and was identified as mixed-layer illite/smectite-1M. Hungarian investigations on Füzérradvány illite in many respects preceded the corresponding American studies but whatever, they are in accordance with them (see: VICZIÁN 1996, 1997).

ERDÉLYI et al. (1957a, b) described a high temperature hydrothermal illite-2M variety which he called “hydro-muscovite”.

From the Mecsek uranium deposits Cr-bearing mica was identified by KISS (1960) and further specified by SZTRÓKAY & PUSKÁS (GÁL-SÓLYMOS et al. 1994).

TOKODY (1962) described a smectite variety as an alteration product of andesite and he called it “mauritzite”. This mineral was later characterized by KÁKAY SZABÓ (1983) and identified as saponite with a high iron content by WEISZBURG et al. (1993).

J. BARNA (1983) described it as a natural — clay-humic substance complex originating from brown coals. It was especially the presumed hydrous varieties in the serpentine group that were extensively studied by ERDÉLYI et al. (1957a, b) and ERDÉLYI & VENIALE (1970).

In a later period PAPP (1993) studied the intergrowth of serpentine minerals and DÓDONY (1997), DÓDONY et al. (2001) published electron microscopic structural analyses of the serpentine varieties antigorite and polygonal serpentine.

The regular mixed-layer mineral “allevardite” (i.e. rectorite) was recognised by NEMECZ et al. (1963) from hydrothermal deposits at Király-hegy in the Tokaj Mts. Another regular mixed-layer clay mineral — corrensite — was first described from Hungary by VICZIÁN (1993); its occurrence was in the German-type Triassic formations of the Mecsek Mts.

### Classification and nomenclature

Immediately before the formation of the Hungarian Clay Group a proposition on classification was submitted jointly by Hungarian specialists for the AIPEA and published by FÖLDVÁRI-VOGL (1958). Already this classification contained the term “couches intercalées” (approximately identical with “interstratified”) and “sárospatakite” was included into this group.

WEISZBURG et al. (2004, 2008) contributed much to the chemical systematics of Fe-bearing members of the mica and smectite groups (e.g. celadonite, glauconite, and with recognition of the “Fe-rich montmorillonite” series).

### Determinative methods in clay mineralogy

A so-called “direct” method of quantitative analysis by X-ray diffraction was developed by NÁRAY-SZABÓ & PÉTER (1967). This was improved later by several authors including BÁRDOSSY et al. (1980), RÍSCHÁK (KOMKOV et al. 1989, SIDORENKO et al. 1992) and SAJÓ (FERET et al. 1997). A special apparatus for thermal analysis — called the “Derivatograph” — was constructed by PAULIK et al. (1986). The apparatus was manufactured in Hungary for several years. An early Atlas of thermal analysis was compiled by FÖLDVÁRI-VOGL (1958).

Later this method was developed to a high standard on the international level by FÖLDVÁRI (1986, 1991). SZÖŐR & BOHÁTKA (1985) combined derivatographic analysis with the mass spectrometry of escaping gases. SZENDREI (2001) applied the analysis of specially prepared thin sections to describe the micro-morphology and genetic processes of soils. The radioactive age of clay substances is determined mainly by the K-Ar method devised by BALOGH (ÁRKAI et al. 1995, 2003), PÉCSKAY et al. (2005, 2006) and their co-workers.

### Soils

STEFANOVITS & DOMBÓVÁRINÉ (1985) published a map of clay minerals in Hungarian soils. Typical Hungarian saline soils were studied by SZENDREI (1985). Red clays, which proved to be mostly relicts or redeposited palaeosoils, were studied by BIDLÓ (1980, 1985), FEKETE & STEFANOVITS (1998) and BERÉNYI ÜVEGES et al. (2003). SCHWEITZER & SZÖŐR (1997) correlated mineralogy and the age of red clays. NEMECZ & CSIKÓS-HARTYÁNI (1995) carried out detailed mineralogical analyses on finely-separated grain fractions of soils.

### Lacustrine clay sediments, bentonites

Sediments from Lake Balaton were studied by a research team led by CSERNY (2002). Specific fossil lake sediments of the craters of basalt volcanoes were recognised by SOLTI and analysed by Zs. BARNA & FÖLDVÁRI (1996). JUHÁSZ (1989) identified the principal component of the basaltic bentonites as Fe-bearing beidellite. It was mainly rhyolite-related bentonite deposits of the Carpathian Basin that were systematically reviewed by KOVÁCS-PÁLFFY (1998). Fine details of the sedimentation conditions of a bentonite deposit were revealed using sequence stratigraphic and geochemical analysis (PÜSPÖK et al. 2005, 2008).

### Lithostratigraphic applications, diagenesis

The clay minerals of Hungarian sedimentary formations were systematically characterised by VICZIÁN (1995). This type of clay study was combined with geochemical research carried out by VARGA et al. (2007), RAUCSIK & VARGA (2008). The diagenetic transformation of smectite into illite was applied for hydrocarbon exploration by VICZIÁN (HÁMOR-VIDÓ & VICZIÁN 1993, TANÁCS & VICZIÁN 1995), J. MÁTYÁS (HILLIER et al. 1995) and others.

### Phyllosilicates in very low grade metamorphic rocks

The very low grade metamorphic stage of transformation has been studied by ÁRKAI et al. (2002). Together with M. TÓTH he improved the “illite crystallinity” (Kübler index) method and introduced the characteristic parameter “chlorite crystallinity” (Árkai index), based on X-ray diffraction (ÁRKAI 1991, ÁRKAI et al. 1996). In addition to potassic white mica paragonite and margarite were also studied (LIVI et al. 2008). The X-ray results were compared with textural analysis using the TEM method.

In cooperation with P. Horváth, Judik etc., his studies extended to the anchimetamorphic formations of the Swiss Alps, also south-eastern territories of Europe (ÁRKAI et al. 1995, JUDIK et al. 2003).

## Hydrothermal alteration of volcanic rocks

The classical studies of PÁLFY (1911) and INKEY (1906) found a correlation between propylitic alteration and gold mineralization in andesitic rocks. Propylitization is a large-scale alteration process in which chlorite forms in place of mafic silicates. SZÁDECZKY-KARDOSS (1958, 1960) drew attention to the uptake of volatiles by magma (transvaporation), the crystallisation of volatile-rich magmas (formation of hypomagmatic rocks) and alteration due to secondary processes (meta-magmatic rocks) which produce various clay minerals. In the Tokaj Mts, zones of *hydrothermal alteration* were first recognised by SZÉKY-FUX (1970) and later intensively studied by NEMECZ et al. (1963), NEMECZ & VARJÚ (1970). In the Velence Hills a deeper level of hydrothermal alteration was explored by the team of DARIDA-TICHY, FÖLDVÁRI, FARKAS, led by I. HORVÁTH (DARIDÁNÉ TICHY et al. 1984). Recently in both regions this line of research has been continued mainly by MOLNÁR (MOLNÁR et al. 1999, PÉCSKAY et al. 2005).

## Economic and environmental geology

A comprehensive inventory of the economic clay deposits of the historical Hungary (i.e. the pre-1920 borders of Hungary) was prepared by KALECSINSZKY (1905). The kaolin, bentonite, illite, zeolite and siliceous earth deposits of the Tokaj Mts and other areas of the country were studied for several decades by E. MÁTYÁS (1966, 1974) and ZELENKA (1994, PÉCSKAY et al. 2005). A comprehensive review was compiled by VÉGHNÉ (1967).

Model experiments of *environmental pollution* by heavy metals were carried out by NÉMETH et al. (2005) and applied to soils by SIPOS (2006–2007). The interaction of high arsenic groundwater with the rock-forming minerals was studied by VARSÁNYI & Ó. KOVÁCS (2005).

Clay liners for communal *waste depositories* were developed by SZABÓ (CZURDA & SZABÓ ed. 1996). Pelitic host rocks and the fault gouge of the granitic host rock for

radioactive waste depositories were studied in Mecsek Mts (FÖLDVÁRI 2006, R. VARGA et al. 2005).

## Extraterrestrial applications

GUCSIK (GAVIN et al. 2010) has started to develop models for the formation of the recently discovered clay minerals on the planet Mars.

## Conclusions

Three main periods can be distinguished in the history of the Hungarian clay mineral research.

In the first period which comprises roughly the 1960–70's, the main interest was focused to the study of hydrothermal alteration in volcanic regions, including the study of specific minerals and economic deposits.

In the second period, approximately in the 1980–90's, investigations concentrated to the study of sedimentary formations, including shales, pelitic rocks and bauxites. Special attention was given to diagenetic processes related to hydrocarbon genesis and very low grade metamorphism.

In the most recent period, on the turn of the 20<sup>th</sup> and 21<sup>st</sup> centuries the importance of young sediments such as soils and Quaternary deposits is growing. There is an increasing interest to environmental applications such as lining of radioactive waste depositories and heavy metal pollution. Thermal analysis was applied on high level in the former and in this period. There are instrumental analyses of specific mineral groups like serpentines and iron clay minerals of high international standard.

In general, the research was able to attain a reliable, medium to good level which matched the international standards. In a few fields, however, such as in the study of bauxites, very low grade metamorphism or in some mineralogical studies Hungarian researchers played a pioneering role in the general clay science.

## References — Irodalom

- ÁRKAI, P. 1991: Chlorite crystallinity: an empirical approach and correlation with the illite crystallinity, coal rank and mineral facies as exemplified by Paleozoic and Mesozoic rocks of northeast Hungary. — *Journal of Metamorphic Geology* **9**, 723–734.
- ÁRKAI, P., BALOGH, K. & DUNKL, I. 1995a: Timing of low-temperature metamorphism and cooling of the Paleozoic and Mesozoic formations of the Bükkium, innermost Western Carpathians, Hungary. — *Geologische Rundschau* **4**, 334–344.
- ÁRKAI, P., SASSI, F. P. & SASSI, R. 1995b: Simultaneous measurements of chlorite and illite crystallinity: a more reliable tool for monitoring low- to very low grade metamorphism in metapelites. A case study from the Southern Alps (NE Italy). — *European Journal of Mineralogy* **7**, 1115–1128.
- ÁRKAI, P., MERRIMAN, R. J., ROBERTS, B., PEACOR, D. R. & TÓTH, M. 1996: Crystallinity, crystallite size and lattice strain of illite-muscovite and chlorite: comparison of XRD and TEM data for diagenetic to epizonal pelites. — *European Journal of Mineralogy* **8/5**, 1119–1137.
- ÁRKAI, P., SASSI, F. P. & DESMONS, J. 2002: Towards a unified nomenclature in metamorphic petrology: 4. Very low- to low grade metamorphic rocks. — A proposal on behalf of the IUGS Subcommission on the Systematics of Metamorphic Rocks. Provisional recommendations, web version of 31.10.2002. (<http://www.bgs.ac.uk/SCMR>), 1–12.
- ÁRKAI, P., FARYAD, S. W., VIDAL, O. & BALOGH, K. 2003: Very low-grade metamorphism of sedimentary rocks of the Meliata unit, Western Carpathians, Slovakia: implications of phyllosilicate characteristics. — *International Journal of Earth Sciences (Geologische Rundschau)* **92**, 68–85.

- BÁRDOSSY, Gy. 1982: *Karst bauxites. Bauxite deposits on carbonate rocks.* — Akadémiai Kiadó, Budapest, 441 p.
- BÁRDOSSY, Gy. & ALEVA, G. J. J. 1990: *Lateritic bauxites.* — 624 p., Elsevier, Amsterdam etc.
- BÁRDOSSY, Gy., BOTTYÁN, L., GADÓ, P., GRIGER, Á. & SASVÁRI, J. (1980): Automated quantitative phase analysis of bauxites. — *American Mineralogist* **65**, 135–141.
- BARNA, J. 1983: Humic substances-clay complexes in Hungarian coals. — *Fuel* **62**, 380–388.
- BARNA, Zs. & FÖLDVÁRI, M. 1996: Bazaltbentonitok termoanalitikai vizsgálata (Thermoanalytical examinations of basaltic bentonites). — In: SOLTI G. (szerk.): Olajpala, alginít, bazaltbentonit kutatása és hasznosítása, 1987–1993 (Prospecting and utilisation of oil shale, alginite, basalt bentonite). 77–80. Alginít Alapítvány, Budapest.
- BERÉNYI ÚVEGES, J., HORVÁTH, Z., MICHLÉI, E., MINDSZENTY, A. & NÉMETH, T. 2003: Reconstructing Quaternary pedogenesis in a paleosol sequence in Hungary. — *Quaternary International* **106–107**, 61–71.
- BIDLÓ, G. 1980: Mineralogical investigation of degraded kaolinites from Dunántúl (Transdanubian) area. — *Acta Mineralogica-Petrographica, Szeged* **24**, Supplementum 111–114. (Proc. 10<sup>th</sup> Kaolin Symp., Budapest, 1979).
- BIDLÓ, G. 1985: Mineralogical investigation of Middle Pliocene and Pliocene–Pleistocene transitional clays. — *5<sup>th</sup> Meeting of the European Clay Groups, Prague, 1983*, 111–115.
- CSENYI T. 2002: A balatoni negyedidőszaki üledékek kutatási eredményei (Results of the investigations on Quaternary sediments of Lake Balaton). — *Földtani Közlöny* **132/különszám** (special issue), 193–213.
- CZURDA, K. & SZABÓ, I. (ed.) 1996: Das Multibarrierensystem in der Deponiebautechnik. Hulladékkelhelyezés többszörös védelmi rendszere. — MBS tanfolyam jegyzet (MBS lecture notes), Budapest, 1996. Schr. Angew. Geol. Karlsruhe **44**.
- DARIDÁNÉ TICHY M., HORVÁTH I., FARKAS L. & FÖLDVÁRI M. 1984: Az andezitmigmatizmushoz kapcsolódó kőzetelváltozások a Velencei-hegység keleti részén (Rock alteration related to andesite magmatism on eastern part of Velence Mts.). — *MÁFI Évi Jelentés (Annual Report of the Geological Institute of Hungary)* **1982-ről**, 271–288.
- DÓDONY, I. 1997: Structure of the 30-sectored polygonal serpentine. A model based on TEM and SAED studies. — *Phys. Chem. Minerals* **24**, 39–49.
- DÓDONY, I., PÓSFAI, M. & BUSECK, P. R. 2001: Revised structure models for antigorite: A HRTEM study. — *American Mineralogist* **87**, 1443–1457.
- ERDÉLYI, J. & VENIALE, F. 1970: Idro-crisotilo: un nuovo minerale del gruppo del serpentino. — *Rend. Soc. Ital. Mineral. Petrol.* **26**, 403–404.
- ERDÉLYI, J., KOBLENČZ, V., TOLNAY, V. 1957a: Einige neue Mineralvorkommen aus der Erzschürfung von Nagybörzsöny (Deutschpilsen). — *Acta Mineralogica-Petrographica, Szeged* **10**, 3–13.
- ERDÉLYI, J., KOBLENČZ, V. & VARGA, N. S. 1957b: Neuere strukturelle Regeln der Hydroglimmer. Hydroantigorit, ein neues Serpentinmineral und metakolloidaler Brucit vom Csódi-Berg bei Dunabogdány (Ungarn). — *Acta Geologica Hungarica* **6**, 95–106.
- FEKETE J. 1988: *Trópusi talajok* (Tropical soils). — Akadémiai Kiadó, Budapest.
- FEKETE, J. & STEFANOVITS, P. 1998: Comparative study of the mineral composition of red clays in Hungary. — *Agrokémia és Talajtan (Agrochemistry and Soil Science)* **47**, 1–4, 23–28.
- FERET, F., AUTHIER-MARTIN, M., SAJO, I. 1997: Quantitative phase analysis of Bidi-Koum bauxites (Guinea). — *Clays and Clay Minerals* **45/3**, 418–427.
- FÖLDVÁRI M. 1986: A földtani kutatásban alkalmazott termoanalitikai módszerek (Thermo-analytical methods applied in geological research). — *MÁFI Módszertani Közlemények (Methodological Contributions, Geol. Inst. of Hung.)* **9/1**, 1–70.
- FÖLDVÁRI, M. 1991: Measurement of different water species in minerals by means of thermal derivatography. — In: SMYKATZ-KLOSS, W. & WARNE, S. St. J. (eds): *Thermal Analysis in the Geosciences*. pp. 84–100. Lecture Notes in Earth Sciences vol. **38**, Springer, Berlin etc.
- FÖLDVÁRI M. 2006: Termoanalízis alkalmazása radioaktív hulladéklerakók földtani kutatása során (Application of thermal analysis for the geological investigation of the disposal of radioactive waste). — *Acta GGM Debrecina, Geology, Geomorphology, Physical Geography Ser.* **1**, 25–32.
- FÖLDVÁRINÉ VOGL M. 1958: *A differenciális termikus elemzés szerepe az ásványtanban és a földtani nyersanyagkutatásban (Differential thermal analysis applied in mineralogy and geological prospecting).* — Műszaki Könyvkiadó, Budapest. 90 p.
- GÁL-SÓLYMOS, K., PUSKÁS, Z., KUBOVICS, I. & VINCZE, J. 1994: Electron microprobe study of the mineral paragenesis in the Mecsek U-ore deposit. — *Acta Mineralogica-Petrographica, Szeged* **35**, Suppl., 13–31.
- GAVIN, P., CHEVRIER, V., NINAGAWA, K., GUCSIK, A. & HASEGAWA, S. 2010: Experimental investigation into the effects of meteoritic impacts on the spectral properties of phyllosilicates on Mars (abstract). — *41st Lunar and Planetary Science Conference, The Woodlands, Texas, U.S.A., 2010*. Abstract No. 1890.
- HÁMOR-VIDÓ, M. & VICZIÁN, I. 1993: Vitrinite reflectance and smectite content of mixed-layer illite/smectites in Neogene sequences of the Pannonian Basin, Hungary. — *Acta Geologica Hungarica* **36/2**, 197–209.
- HILLIER, S., MÁTYÁS, J., MATTER, A. & VASSEUR, G. 1995: Illite/smectite diagenesis and its variable correlation with vitrinite reflectance in the Pannonian Basin. — *Clays and Clay Minerals* **43/2**, 174–183.
- INKEY, B. 1906: De la relation entre l'état propylitique des roches andesitiques et leurs filons minéraux. — *C. R. Congr. Géol. Internat., Xème session, Mexico, 1906*, **1**, 501–517.
- JUDIK, K., NÉMETH, T., TIBLJAŠ, D., HORVÁTH, P. & ÁRKAI, P. 2003: Corrensite in metabasalts and metagabbros from Mt. Medvednica, Croatia. — *Acta Mineralogica-Petrographica, Szeged* **44**, 1–8.
- JUHÁSZ A. Z. & OPOCZKY L. 1982: *Szilikátok mechanikai aktiválása finomőrléssel.* — Szilikátkémiai Monográfiák XI. Akadémiai Kiadó, Budapest. 244 p.
- JUHÁSZ, A. Z. & OPOCZKY, L. 1990: *Mechanical activation of minerals by grinding: puverizing and morphology of particles.* — Ellis Horwood, Chichester, England. 234 p.

- JUHÁSZ A. Z. 1995–96: Általános és szilikátkémiai kolloidika (General and silicate colloid chemistry). **1** (1995), 200 p. **2** (1996), 165 p., **3** (1996), 200 p.—Egyetemi jegyzet (university course), Veszprémi Egyetem, Veszprém.
- JUHÁSZ Z. 1989: A várkeszői bentonittípus technológiai tulajdonságai (The technological properties of the bentonite type of Várkesző).—*Földtani Kutatás* **32/4**, 65–70.
- KÁKAY SZABÓ O. 1983: A mauritzit újrizsgálata (Re-examination of mauritzite).—*Földtani Közlöny* **113/4**, 333–356.
- KALECSINSZKY S. 1905: A magyar korona országainak megvizsgált agyagai (Clays examined in the lands of the Hungarian Crown).—A M. Kir. Földtani Intézet kiadványa (Published by the Hung. Royal Geol. Inst.), Budapest. 218 p., 1 térképmelléklet.
- KISS J. 1960: Az urán–króm–vanadium eloszlása és az epigén krómcsillám szerepe a mécsei permie összetben (Die Verteilung von U–Cr – V und die Rolle des epigenetischen Chromglimmers im Permkomplex des Mecsekgebirges).—*Földtani Közlöny* **90/1**, 73–82.
- KOMKOV, A. I., DYAKONOV, YU. S., MISCHENKO, K. S., RAYNOV, N., CECHLAROVA, I., RISCHÁK, G., UNGER, H. & HERING, A. 1989: Application of quantitative X-ray diffraction phase analysis in the geological survey. I. A methodological guide. — *Scientific Commission of Analytical Methods, Scientific Commission of Methods of Mineralogical Research, VIMS, Moskva* (in Russian).
- KOVÁCS-PÁLFFY P. 1998: Harmadidőszaki bentonit típusú ásványi nyersanyagtelepek ásványtani, geokémiai és genetikai összehasonlító vizsgálata (Comparative mineralogical, geochemical and genetic study of Tertiary bentonite-type deposits). — *Ph. D. értekezés összefoglaló tézisei* (Ph. D. Thesis). Kossuth Lajos Tudományegyetem, Debrecen.
- LIVI, K. J. T., CHRISTIDIS, G. E., ÁRKAI, P. & VEBLEN, D. R. 2008: White mica domain formation: A model for paragonite, margarite, and muscovite formation during prograde metamorphism. — *American Mineralogist* **93/4**, 520–527.
- M. NAGY, N. & KÓNYA, J. 2009: Interfacial chemistry of rocks and soils. — *Surfactant Science Series*, **148**, CRC Press, Boca Raton, FL, U.S.A. 244 p.
- MAEGDEFRAU, E. & HOFMANN, U. 1937: Glimmerartige Mineralien als Tonsubstanzen. — *Zeitschrift für Kristallographie* **98/1**, 31–59.
- MÁTYÁS E. 1966: A rátkai felsőszarmata édesvízi medence földtani és teleptani viszonyai (General and economic geological relations of the Rátka Upper Sarmatian limnic basin).—*Földtani Közlöny* **96/1**, 27–42.
- MÁTYÁS, E. 1974: Volcanic and postvolcanic processes in the Tokaj Mountains on the basis of geological data of raw material prospecting. — *Acta Geologica Hungarica* **18/3–4**, 421–455.
- MINDSZENTY A. 1999: Bauxitszedimentológia. A karsztauxitok, a geodinamika és a klíma kapcsolatának elemzése mediterrán példákban (Bauxite sedimentology. Analysis of the relationship between karst bauxites, geodynamics and climate on Mediterranean examples). — *Akadémiai doktori értekezés* (D.Sc. Thesis), Budapest. 1–171.
- MOLNÁR, F., LEXA, J. & HEDENQUIST, J. W. 1999: Epithermal mineralization of the Western Carpathians. — Guidebook Prepared for the Society of Economic Geologists Field Conference, 4–7 September 1999, Society of Economic Geologists Guidebook Series **31**.
- NÁRAY-SZABÓ, I. & PÉTER, É. 1967: Die quantitative Phasenanalyse in der Tonmineralforschung. — *Acta Geologica Hungarica* **11/4**, 347–356.
- NEMECZ, E. 1981: *Clay minerals*. — Akadémiai Kiadó, Budapest, 547 p.
- NEMECZ E. 2006: Ásványok átalakulási folyamatai talajokban (Transformation processes of minerals in soils). — Akadémiai Kiadó, Budapest. 324 p.
- NEMECZ, E. & CSIKÓS-HARTYÁNI, Zs. 1995: Processes in soils and paleosoils. — *GeoJournal* **36/2–3**, 139–142.
- NEMECZ, E., VARJU, Gy. & BARNA, J. 1963: *Allevardite from Királyhegy, Tokaj Mountains, Hungary*. — *Proc. Intern. Clay Conf. Stockholm* **2**, 51–67. Pergamon Press, Oxford etc.
- NEMECZ, E. & VARJU, Gy. 1970: Sárospatakít (Hidrotermális illit-montmorillonit) kémiái és szerkezeti sajátságai (Chemical and structural investigation of Sárospatakites, Illite/montmorillonite). — *Földtani Közlöny* **100/1**, 11–22. (in Hungarian)
- NÉMETH, T., MOHAI, I. & TÓTH, M. 2005: Adsorption of copper and zinc ions on various montmorillonites: an XRD study. — *Acta Mineralogica-Petrographica, Szeged* **46**, 29–36.
- PÁLFY M. 1911: Az Erdélyrészki Érchegység bányáinak földtani viszonyai és érctelérei (Die geologischen Verhältnisse und die Erzlagerstätten des siebenbürgischen Erzgebirges). — *A Magyar Királyi Földtani Intézet Évkönyve (Annals of the Royal Geological Institute of Hungary)* **18/4**, 205–464.
- PAPP, G. 1993: Oriented intergrowth of crysotile with lizardite and antigorite. — *Neues Jahrbuch für Mineralogie Monatshefte* **1993/1**, 1–9.
- PAULIK, J., PAULIK, F. & ARNOLD, M. 1986: Derivatograph-c. A microcomputer automated equipment for simultaneous TG, DTG, DTA, EGA and TD. — *Thermochimica Acta* **107**, 375–378.
- PÉCSKAY, Z., MOLNÁR, F., ITAYA, T., ZELENKA, T. 2005: Geology and geochronology of illite from the clay deposit at Füzérradvány, Tokaj Mts., Hungary. — *Acta Mineralogica-Petrographica, Szeged* **46**, 1–7.
- PÉCSKAY Z., MOLNÁR, F., ITAYA, T., ZELENKA, T., SEGHEDI I., BALOGH K., KONECNY V., ZELENKA T., KOVÁCS M., PÓKA T., FÜLÖP A., MÁRTON E., PANAIOTU C. & CVETKOVIC V. 2006: Geochronology of Neogene Magmatism in the Carpathian arc and intra-Carpathian area. — *Geologica Carpathica* **57/6**, 511–530.
- PÜSPÖKI, Z., KOZÁK, M., KOVÁCS-PÁLFFY, P., FÖLDVÁRI, M., MCINTOSH, R. W. & VINCZE, L. 2005: Eustatic and tectonic/volcanic control in sedimentary bentonite formation – a case study of Miocene bentonite deposits from the Pannonian Basin. — *Clays and Clay Minerals* **53/1**, 71–91.
- PÜSPÖKI, Z., KOZÁK, M., KOVÁCS-PÁLFFY, P., SZEPESI, J., MCINTOSH, R., KÓNYA, P., VINCZE, L. & GYULA, G. 2008: Geochemical records of a bentonitic acid-tuff succession related to a transgressive systems tract – indication of changes in the volcanic sedimentation rate. — *Clays and Clay Minerals* **56/1**, 23–38.
- R. VARGA, A., SZAKMÁNY, Gy., RAUCSIK, B. & MÁTHÉ, Z. 2005: Chemical composition, provenance and early diagenetic processes of playa lake deposits from the Boda Siltstone Formation (Upper Permian), SW Hungary. — *Acta Geologica Hungarica* **48/1**, 49–68.
- RAUCSIK B. & VARGA A. 2008: Az alsó-toarci feketepala Réka-völgyi szelvényének ásványtani jellemzése (Óbányai Aleurolit Formáció, Mecsek hegység): őséghajlattani következtetések (Mineralogy of the Lower Toarcian black shale section from the Réka Valley (Óbányai Siltstone Formation, Mecsek Mountains, Hungary): implications for palaeoclimate). — *Földtani Közlöny* **138/2**, 133–146.

- SCHWEITZER, F. & SZÖÖR, Gy. 1997: Geomorphological and stratigraphic significance of Pliocene red clay in Hungary. — *Zeitschrift für Geomorphologie N.F. Suppl.-Bd.* **110**, 95–105.
- SIDORENKO, G. A., VOLKOV, M. A., DYAKONOV, Yu. S., MISCHENKO, K. S., RAYNOV, N., CECHLAROVA, I., RISCHÁK, G., MELKA, K., KORECKY, J., UNGER, H. & HERING, A. 1992: Application of quantitative X-ray diffraction phase analysis in the geological survey. II. 1–28. A methodological guide. — Scientific Comission of Methods of Mineralogical Research, VIMS, Moskva (in Russian).
- SIPOS, P. 2006–2007: Mobilization conditions of lead in forest soils from the Cserhát Mts., NE Hungary. — *Acta Mineralogica-Petrographica, Szeged* **47**, 53–59.
- STEFANOVITS P. 1981: *Talajtan (Soil science)*. — 2. átdolg. kiadás. Mezőgazdasági Kiadó, Budapest.
- STEFANOVITS P. & DOMBÓVÁRI L-NÉ (K.) 1985: A talajok agyagásvány-társulásainak térképe (The map of clay mineral associations in the soils of Hungary). — *Agrokémia és Talajtan* **34/3–4**, 317–330.
- SZÁDECZKY-KARDOSS, E. 1958: On the petrology of volcanic rocks and the interaction of magma and water. — *Acta Geologica Hungarica* **5**, 197–233.
- SZÁDECZKY-KARDOSS, E., PANTÓ, G. & SZÉKY-FUX, V. 1960: A preliminary proposition for developing a uniform nomenclature of igneous rocks. — *International Geological Congress, Report of the 21<sup>st</sup> Session Norden, 1960, Part 13*, 287–292.
- SZÁNTÓ F. 1987: *A kolloidkémia alapjai (Basic colloid chemistry)*. — Gondolat, Budapest. 337 p.
- SZÉKYNÉ FUX V. 1970: *Telkibánya ércessedése és kárpáti kapcsolatai (The Telkibánya mineralisation and its Carpathian relations)*. — Akadémiai Kiadó, Budapest. 266 p.
- SZENDREI G. 1979: A magyar agyagásványtani szakirodalom jegyzéke, 1900–1978 (Bibliography of Hungarian clay mineralogy, 1900–1978). — *Kézirat (Manuscript)*, Budapest.
- SZENDREI, G. 1985: The stability and distribution of clay minerals in Hungarian salt affected soils. — *5th Meeting of the European Clay Groups, Prague, 1983*. 471–476. Charles University, Prague.
- SZENDREI G. 1994: Talajásványtan (Soil mineralogy). — *MÁFI Módszertani Közlemények (Methodological Contributions, Geol. Inst. of Hung.)* **1994/1**, 1–217.
- SZENDREI G. 2001: *A hazai talajtípusok mikromorfológiája (Micromorphology of soil types in Hungary)*. — A szerző kiadása (Edition of the author), Budapest. 163 p.
- SZENDREI G. (compiled by) 2010: A Szakosztály tisztségsviselői által publikált monografiák, könyvek (Monographs, books published by the officers of the Clay Group). — *Kézirat (Manuscript)*, Budapest.
- SZÖÖR, Gy. & BOHÁTKA, S. 1985: Derivatograph-Qms system in geochemical research. — *Thermochimica Acta* **92**, 395–398.
- TANÁCS, J. & VICZIÁN, I. 1995: Mixed-layer illite/smectites and clay sedimentation in the Neogene of the Pannonian Basin, Hungary. — *Geologica Carpathica, ser. Clays* **4/1**, 3–22.
- TOKODY, L. 1962: Mauritzite ein selbständiges Mineral. — *Annales hist.-nat. Musei Nationalis Hung., Pars Min. Pal.* **54**, 27–30.
- VARGA, A., SZAKMÁNY, Gy., ÁRGYELÁN, T., JÓZSA, S., RAUCSIK B. & MÁTHÉ Z. 2007: Complex examination of the Upper Paleozoic siliciclastic rocks from southern Transdanubia, SW Hungary – mineralogical, petrographic, and geochemical study. — In: ARRIBAS, J., CRITELLI, S. & JOHNSSON, M. J. (eds): Sedimentary Provenance and Petrogenesis. *Geological Society of America, Special Paper* **420**, 221–240.
- VARSÁNYI, I. & Ó. Kovács, L. 2005: The role of groundwater flow in controlling the arsenic concentration in the southern part of the Great Hungarian Plain. — *Acta Mineralogica-Petrographica, Szeged* **46**, 47–52.
- VÉGH S-NÉ 1967: *Nemércek földtana (Geology of non-metallic mineral deposits)*. — Tankönyvkiadó, Budapest. 283 p.
- VICZIÁN, I. 1993: Clay mineralogy of Middle Triassic evaporitic and carbonate rocks, Mecsek Mts. (southern Hungary). — *11th Conference on Clay Mineralogy and Petrology, Č. Budějovice, 1990*, 135–144. Univerzita Karlova, Praha.
- VICZIÁN, I. 1995: Clay minerals in Mesozoic and Paleogene sedimentary rocks of Hungary. — *Romanian Journal of Mineralogy* **77**, 35–44.
- VICZIÁN, I. 1996: A füzérradványi illit vizsgálatának újabb eredményei (New results in the study of the Füzérradvány illite). — *Földtani Közlöny* **126/2–3**, 263–266.
- VICZIÁN, I. 1997: Hungarian investigations on the “Zempleni” illite. — *Clays and Clay Minerals* **45/1**, 114–115.
- VICZIÁN, I. 1999: Brief historical notes: The Hungarian Clay Mineral Group. — *EGCA Newsletter* **3**, 1999, 22–23.
- VICZIÁN, I. 2000: History of mineralogical investigations of the Füzérradvány „illite”, near Sárospatak, Hungary. — *Acta Geologica Hungarica* **43/4**, 493–500.
- VOGL M. 1982: Az agyagásványok vizsgálati módszereinek hazai fejlődése az elmúlt 20 év alatt (Advance of the analytical methods in clay mineralogy during the last 20 years in Hungary). — *Földtani Közlöny* **112/1**, 31–39.
- WEISZBURG, T., BADA, G., DÓDONY, I., JÁNOSI, M., LOVAS, Gy., NAGY-BALOGH, J., CZAKÓ-NAGY, I., NAGY, S. & PAPP, G. 1993: A „mauritzit”, Erdőbénye nevezetes szaponit-változata („Mauritzite”, a special saponite variety from Erdőbénye, NE Hungary). — *Topographia Mineralogica Hungariae* **1**, 57–67. Herman Ottó Múzeum, Miskolc.
- WEISZBURG, T. G., TÓTH, E. & BERAN, A. 2004: Celadonite, the 10-Ågreen clay mineral of the manganese carbonate ore, Úrkút, Hungary. — *Acta Mineralogica-Petrographica, Szeged* **45/1**, 65–80.
- WEISZBURG, T. G., TÓTH, E., POP, D. & CORA, I. 2008: Iron dominated dioctahedral TOT clay minerals: from nomenclature to formation processes (abstract). — *MECC'08, Zakopane, 2008, Abstracts. Mineralogia, Spec. Papers* **33**, 31.
- ZELENKA, T. 1994: Genetic relationship of the Hungarian clay deposits (Kaolin-bentonite). — *Acta Mineralogica-Petrographica, Szeged* **25**, 97–101.