

Spatial analysis of changes and anomalies of intense rainfalls in Hungary

GERGELY JAKAB^{1,2,6}, TIBOR BÍRÓ³, ZOLTÁN KOVÁCS⁴, ÁDÁM PAPP⁴, NINSAWAT SARAWUT⁵,
ZOLTÁN SZALAI^{1,6}, BALÁZS MADARÁSZ¹ and SZILÁRD SZABÓ⁴

Abstract

Extreme precipitation events can trigger flash flood, mass movements, pluvial flood and accelerated soil erosion. As soil structures are highly degraded due to intensive improper cultivation water infiltration can considerably decrease during the vegetation period. Additional changes in canopy coverage on the soil surface cause relevant variability in infiltration and hence vulnerability against runoff related disasters. Most researchers agree that the frequency of extreme precipitations increases, however, in the Carpathian Basin the uncertainties are quite high. This study aims to compare daily maximum mean precipitation amounts (MMPA) predicted by the Goda-method for June and August as the most probable months of extremities. We used the CarpatClim database as input and predicted MMPAs for two periods, 1960–1985 and 1986–2010. The Goda-method uses monthly data and calculates daily results on given probability. A general increase was found between the first and second half of the period regarding daily maximum precipitation amount in both investigated months. For August the 1-day precipitation amount increased from 56.1 mm to 61.8 mm, whereas 6-days amount from 93.8 mm to 103.2 mm at 1 per cent probability ($r = 0.53$; $p < 0.001$). Beyond this change, relevant spatial differences were found. Comparing the macro regions plains had lower increase compared to the mountains, whereas the highest increase was in the Transdanubian Hills (TH). The most endangered location is the southern part of the Transdanubian Hills where parallel with the intensive increase in MMPA both in June and August the environmental conditions such as loose parent material and the high percentage of crop fields also emphasize the potential hazard.

Keywords: extreme precipitation, climate change, soil erosion, flash flood, pluvial flood

Introduction

Climate change can trigger harmful effects in urban areas as well (LÁSZLÓ, E. *et al.* 2016), however, the most dangerous changes are predicted for agricultural areas. A well-developed soil can take and keep even 200–300 mm precipitation at one time depending on its land-use, porosity, aggregation and canopy cover. However, soils especially those

of crop fields and vineyards are highly degraded; therefore, water infiltration is inhibited (JAKAB, G. *et al.* 2017; RODRIGO-COMINO, J. 2019). Moreover, since the beginning of the last century, extreme rainfalls occurred more frequently in all over Europe, as a consequence of climate change (FOWLER, H.J. and KILSBY, C.G. 2003; MÜLLER, M. *et al.* 2009; MILOSEVIC, D. and SAVIC, S. 2013; LAKATOS, M. and HOFFMANN, L. 2018).

¹Geographical Institute, Research Centre for Astronomy and Earth Sciences, Hungarian Academy of Sciences, Budapest, Hungary. E-mails: jakab.gergely@csfk.mta.hu, szalai.zoltan@csfk.mta.hu, madarasz.balazs@csfk.mta.hu

²Institute of Geography and Geoinformatics, University of Miskolc, Miskolc, Hungary.

³Faculty of Water Sciences, Institute of Hydraulic Engineering and Water Management, National University of Public Service, Baja, Hungary. E-mail: biro.tibor@uni-nke.hu

⁴Department of Physical Geography and Geoinformatics, Faculty of Science and Technology, University of Debrecen, Debrecen, Hungary. E-mail: szabo.szilard@science.unideb.hu

⁵Department of Remote Sensing and Geographic Information Systems (RS-GIS), Asian Institute of Technology, Bangkok, Thailand. E-mail: sarawutn@ait.ac.th

⁶Department of Environmental and Landscape Geography, ELTE University, Budapest, Hungary.

Parallel with the increasing frequency of extreme precipitation events, rainfall occurrence is getting more erratic (BARTHOLY, J. and PONGRÁCZ, R. 2007; MIKA, J. and FARKAS, A. 2017; MILOSEVIC, D. et al. 2017). An increasing ratio of annual precipitation will not be able to infiltrate, triggering water deficiency during the growing season in a mesic environment (KNAPP, A.K. et al. 2008). The water remaining on the soil surface causes runoff and flash flood on the hillslopes (CZIGÁNY, Sz. et al. 2010), and pluvial floods on lowlands. Both phenomena have primary importance in soil degradation in Hungary as 49 per cent of the territory is arable field and 35 per cent of that is already eroded (KERTÉSZ, Á. and CENTERI, Cs. 2006). Pluvial floods including melting snow and intense rains frequently cover considerable parts of the plains in spring (VAN LEEUWEN, B. et al. 2013; PÁSZTOR, L. et al. 2015; BALÁZS, B. et al. 2018). Summer rainfall events lead to pluvial floods as well for several days in high quantity. To avoid or minimize damages, actual water infiltration should be increased by tillage and plant coverage. As soil infiltration capacity varies in a wide range both in time and in space, agricultural technology should adapt to it in order to fight against runoff and pluvial floods (JAKAB, G. et al. 2017).

The ability of a precipitation event to destroy the soil surface and to generate runoff is called erosivity, which is proportional with the amount and intensity of the storm (WALTNER, I. et al. 2018). Rainfall erosivity (R factor) is generally calculated from these two properties according to the USLE (WISCHMEIER, W.H. and SMITH, D.D. 1978) or the RUSLE (RENARD, K.G. and FREIMUND, J.R. 1994) model. Traditionally, R factor values are determined for a whole calendar year in order to estimate the erosion potential of the precipitation (PANAGOS, P. et al. 2015). Conversely, long-term erosion monitoring has proven that the main part of soil loss is triggered by only one or two extreme precipitation events those also take the main part of the R factor. Since soil porosity and canopy cover varies in a wide range during the vegetation period the actual runoff and

soil loss volume is the function of the date of a given heavy rainstorm. Hence annual R factor needs to be divided into shorter temporal sections such as months (BALLABIO, C. et al. 2017), single precipitation events or even hours (KENDON, E. et al. 2014).

As the occurrence of extreme precipitation events varies in a wide range both in time and space their proper forecast is still an unsolved question (MAHERAS, P. et al. 2018) hence the Carpathian Basin is presumed to be the most questionable area in Europe regarding future trends (KRISTÓF, E. et al. 2017; GELYBÓ, Gy. et al. 2018; LÁSZLÓ, E. and SALAVEC, P. 2018), but scientists agree that the number of intense precipitation events is expected to increase (BARTHOLY, J. and PONGRÁCZ, R. 2007; CHEVAL, S. et al. 2016), the return time of extreme events increases by 1.2–2 times (PIECKA, I. et al. 2011; PONGRÁCZ, R. et al. 2014). Although there is a long-term precipitation database for whole Europe, locally low spatial resolution and/or the lack of (high resolution) rainfall intensity data cause difficulties in extreme precipitation prognostication (ZHANG, F. et al. 2006). Consequently, models were developed to estimate heavy storm frequency at local scales. Some of these models need detailed measured input and calibration such as ALADIN/AROME (IHÁSZ, I. et al. 2018), whereas simpler models use only long-term monthly (RENARD, K.G. and FREIMUND, J.R. 1994) or daily (GODA, L. 1966) precipitation volume data. The Goda-method uses long-term observations, monthly data, but the output is the daily maximum by months at a given probability level; thus, it has the advantage to describe erosivity or the tendency for pluvial floods in a simple way.

Rapid and heavy rainstorms have direct connections with erosion and could be represented as 1-day maximums. Rainfall maximums (the longest statistically evaluable term is 6 days in Hungary according to GODA, L. 1966) refer to pluvial flood formation in the plains caused by the low infiltration capacity and the intense rains where the runoff is limited because of the flat surface. In Hungary, precipitation maximums are in June (1–6 days maximums) and August when mainly extreme

1-day rainfalls occur, hence these months are presumed to have the most important role in soil erosion and partly in pluvial flood formation (SZÚCS, P. et al. 2006; JAKAB, G. et al. 2015).

The aim of the present study is to quantify changes in maximum daily precipitation volumes at the time of the annual precipitation peaks (in June and August) in Hungary. Our hypotheses to be tested are i.) maximum daily precipitation maximums are in strong correlation with elevation; ii) the maximum daily precipitation volume has increased since 1960 in Hungary; iii.) the spatial distribution of the average change has a high standard deviation; iv.) there are significant differences among the macro regions regarding the mean maximum precipitation amount (MMPA) changes.

Methods

Datasets

We used the precipitation data of the CarpatClim (CC) database (SZALAI, S. et al. 2013; SPINONI, J. et al. 2015). CC is a climatic dataset which is the result of the contribution of nine countries. We filtered out the Hungarian data points, however, the database did not

cover the westernmost part of the country. The dataset is a homogenized time series of 50 years from 1960 to 2010 with the Multiple Analysis of Series for Homogenization (MASH) method, based on measured data but the final product is an interpolated 10 km x 10 km grid network with the Meteorological Interpolation based on the Surface Homogenized Data Basis (MISH) method (SZENTIMREY, T. and BIHARI, Z. 2007; SZENTIMREY, T. 2011; SZENTIMREY, T. et al. 2012).

Calculation of intense rainfalls

We applied the method of GODA, L. (1966) to determine the intensive rainfalls. The method uses monthly data and calculates daily results on a given probability. Firstly, we had to determine whether the time series follows a Pearson-3 or Gamma-2 distribution, and then, by applying the univariate statistics, the constants of the chosen distributions and probability level, and the spatially predefined parameter of the region and applying the appropriate function, we get the final result regarding 1 and 6 days (Figure 1). The process of the workflow is described after GODA, L. (1966).




Fig. 1. Workflow of data management. – Cv = coefficient of variation; Cs = coefficient of asymmetry. Pearson 3 and Gamma 2 are distribution types.

Firstly, we determine the mean (\bar{x}) and standard deviation (σ) of the time series, then the next is the calculation of the coefficient of variation (c_v , Eq. 1) and asymmetry (c_s , Eq. 2).

$$c_v = \frac{\sigma}{\bar{x}}, \tag{1}$$

where c_v is the coefficient of variation, σ is the standard deviation, \bar{x} is the mean of the time series.

$$c_s = \frac{\sum_{i=1}^N (k_i - 1)^3}{(N - 1)c_v^3}, \tag{2}$$

where c_s is the coefficient of asymmetry, k_i is the ratio of x_i and \bar{x} , x_i is the i^{th} element, c_v is the coefficient of variation of the time series.

If $\frac{c_s}{c_v} \geq 2$ then the time series follows a Pearson III. (Foster-Ribkin variation) distribution, and if $\frac{c_s}{c_v} < 2$ the distribution can be described with the Gamma II. (Krickij-Menkely variation) distribution.

GODA, L. (1966) provided a formula for the T = 1–6 days mean precipitation maximums (Eq. 3):

$$\bar{x}^{(1)} = \varphi^{(1)} \bar{x}, \tag{3}$$

where $\bar{x}^{(1)}$ is the mean precipitation maximum of 1 day, for $\varphi^{(1)}$ see Eq. 5; the general formula for T = 1–6 days is (Eq. 4):

$$\bar{x}^{(i)} = \varphi^{(i)} \bar{x}, \tag{4}$$

where $\bar{x}^{(i)}$ is the mean of T = i days-long precipitation maximums, $\varphi^{(i)}$ is the spatial factor with the following formula (Eq. 5):

$$\varphi^{(i)} = \frac{\bar{x}^{(1)}}{\bar{x}} + \Delta\varphi^{(i)}, \tag{5}$$

where $\varphi^{(i)}$ is a factor describing the means of i-days-long (T = 1–6) precipitation maximums, $\Delta\varphi^{(i)}$ is the change for the precipitation maximum as a function of the number of days.

The final step is the calculation of the precipitation maximums on a given probability level. If the c_s/c_v is larger than 2 we use the Φ -value from the Foster-Ribkin table, if it is smaller, we use the K-value from the Krickij-Menkely table with the formulas of Eq. 6 and 7:

$$c_p^{(i)} = (\Phi c_v + 1)\bar{x}^{(i)}, \tag{6}$$

where $c_p^{(i)}$ is the mean precipitation maximum on a given probability level, Φ can be determined from the Foster-Ribkin table using the c_s and probability (e.g. 1% or 5%) values.

$$c_p^{(i)} = K\bar{x}^{(i)}, \tag{7}$$

where $c_p^{(i)}$ is the mean precipitation maximum on a given probability level, K can be determined from the Krickij-Menkely table using the c_v and probability (e.g. 1% or 5%).

The procedure was implemented in R 3.5.1 (R Core Team, 2017) environment with the tidyR (WICKHAM, H. and HENRY, L. 2018) package, and the calculations were completely automated.

Statistical analysis

We divided the 50-year dataset into two periods: 1960–1985 and 1986–2010; thus, we were able to identify whether there is a trend in this time range considering the maximums of the intense rainfalls. Even though MIKA, J. and FARKAS, A. (2017) have pointed out, that many weather extremities have already occurred in the 1960s.

From various possible outcomes, we choose the following combinations: 1- and 6-day maximums with 1 per cent and 5 per cent probability regarding June and August. We checked the variables for normal distribution with the Shapiro-Wilk test and, as some variables were skewed, we applied the non-parametric Wilcoxon-test to reveal if there was a significant difference between the two time periods.

We also determined the effect sizes for the comparisons, to get a standardized and comparable value of the magnitude of differences. Effect size ranges from 0 to 1, and 0 means complete similarity and the increase of the value indicates a larger difference: 0.1 denotes weak, 0.3 moderate while 0.5 large difference (FIELD, A. 2009; SULLIVAN, G.M. and FEINN, R. 2012).

For spatial comparison purposes, we applied the macro regions of Hungary (DÖVÉNYI, Z. 2010), even though this classification was landscape based, resulting units with heterogeneous climatic conditions (Figure 2).

Comparisons were conducted between each macro region pairs to highlight the spatial differences in MMPA change between the two investigated periods (2 x 25 years). On one hand, macro regions reflect the spatial character of changes (absolute situation), on the other hand, all macro regions have different average surface height. Thus, a hypothesis testing, carried out with the robust ANOVA test, was able to reveal whether these differences were significant and characteristic in a region, or they happened just by chance having a random distribution. The results were presented in a visual form on the diagram: where the mean and 95 per cent confidence intervals coincide with the zero value, the difference was not considered to be significant. We conducted correlation analysis between elevation and the different representations of the intense rainfalls,

using Spearman rank correlation. Terrain height was derived from SRTM data at each pixel centre of the CC database. Statistical analyses were conducted with R 3.5.1 (R Core Team, 2018) with the coin (HOTHORN, T. et al. 2008) and the walrus (LOVE, J. and MAIR, P. 2017) packages.

Results

Intense rainfall maximum changes on country level

Daily MMPAs during the investigated fifty years changed between 40 and 137 mm depending on time and spatial location (Figure 3). The precipitation range was the same for the whole country in June and August, whereas the maxima were found at four individual locations in June (Figure 3, a) and in central Transdanubia in August (Figure 3, b). The less extreme daily rainfall amounts were in Kisalföld and in North Hungary.




Fig. 2. Macro regions and other spatial units appearing in the study

Correlation analysis revealed that the 1- and 6-days as well as the 1 per cent and 5 per cent probability intense rainfall maximums were strongly correlated, but the correlation between June and August was weak. Accordingly, we focused on the monthly differences, but also analysed the 6-day events because of the higher values.

Based on 50-year data a general increase was found between the first and second half of the period regarding daily maximum precipitation amount in both investigated months. However, the difference was not significant in any case in June on a country level (Table 1).

Taking a more detailed spatial resolution into account it is clear that locally significant differences were found in 6-days' MMPA in June (Figure 4). In central Hungary, the 6-day

MMPA decreases, whereas the highest decrease was in the TH. The increase is located in the western part of the country, in the Northern part of GHP and in the South near Szeged. Conversely, in August the increase is clear and significant concerning both duration and probability. Beyond this general trend again explicit spatial differences exist (Figure 5).

On the major part of the country the extremity decreased, in the central part of the GHP and in Transdanubia the average dropped more than 20 mm. In contrast, in the southern part of Transdanubia, the rainfall amount increased more than 10 mm, which corresponded to a 5–25 per cent rise (compare to Figure 3, b). Therefore, it is not enough to compare the country means, for the identification of the most endangered locations a higher spatial resolution is needed.




Fig. 3. Spatial distribution of daily MMPAs in June (a) and August (b) derived from the CC using the Goda-method

Table 1. Mean and standard variation of the intense rainfall data (mm) by the number of rainy days, months and probabilities in the studied periods

Variation	Day	Month	1960–1985	1986–2010	p*	Effect size (r)
Mean ± SD 1%	1	June	56.5 ± 6.7	57.6 ± 9.1	<0.001	0.13
	1	August	56.1 ± 10.6	61.8 ± 10.1	<0.001	0.53
	6	June	99.1 ± 10.4	100.9 ± 13.6	<0.001	0.12
	6	August	93.8 ± 16.9	103.2 ± 15.7	<0.001	0.53
Mean ± SD 5%	1	June	46.5 ± 5.2	46.8 ± 7.1	0.33	0.03
	1	August	45.1 ± 8.4	49.0 ± 7.6	<0.001	0.52
	6	June	81.6 ± 7.7	81.9 ± 10.6	0.52	0.02
	6	August	75.5 ± 13.5	81.9 ± 11.8	<0.001	0.52

*p = significance of the pairwise comparison with the Wilcoxon test; italics highlights = p < 0.01.




Fig. 4. Change in average 6-days MPPA in June. Differences between averages of 1960–1985 and 1986–2010 in mm at 1 per cent probability level.




Fig. 5. Change in MPPA in August. Differences between averages of 1960–1985 and 1986–2010 in mm at 1 per cent probability level.


Areal comparison of differences of intense rainfall maximums by macro-regions

In June, regarding the 1-day MMPA, the increase in TH was a significantly higher increase than in the other macro region except TM, which was a neighbouring region. In contrast, the increase was significantly lower in KA than in the other macro region except GHP (Figure 6, above). Thus, the highest difference was found between TH and KA (~ 8 mm), which was more than 13 per cent of the KA value (~ 60 mm). As the calculation is derived from the same database the


differences between 6-days long MMPA showed the same pattern with higher values (Figure 6, below).

In August, the largest difference was higher (~15 mm) in the 1-day MMPA case, which was almost 25 per cent difference compared to the KA basis value of KA. This difference was found between TM and TH. Moreover, TH had a significantly higher increase than any other macro region, whereas TM had a lower increase than any other macro regions, even though the difference between TM and KA was not significant.


Correlation analysis revealed significant correlations between elevation and different




Differences of mean rainfall maximums (1% 1-day, June)



Differences of mean rainfall maximums (1% 1-day, August)



Differences of mean rainfall maximums (1% 6-day, June)



Differences of mean rainfall maximums (1% 6-day, August)

Fig. 6. Mean differences (mm) of 1-day and 6-day MMPA at 1 % probability level in June and August by macro-regions (95% confidence intervals coinciding with 0 are not significant differences, $p > 0.05$). GHP = Great Hungarian Plain; KA = Kisalföld (Little Plain); TH = Transdanubian Hills; TM = Transdanubian Mountains; NHM = North Hungarian Mountains.

intense rainfall measures (Figure 7). The best correlations were found in the case of August regardless of the probability level (1% or 5%); nevertheless, the largest value belonged to a 1 per cent representation (but it was only 0.03 larger than the next one which was a 5% representation).

Generally, correlations were > 0.5 in the case of the period of 1960–1985 for 6-day events of Augusts, but for 1-day events in June for the same period, *r* values were below 0.3. For the period of 1986–2010, correlations ranged between 0.32–0.46, and the largest value belonged to the 6-day events in August on 5 per cent probability. However, we did not detect relevant correlation considering the changes between the intense rainfalls and elevation regarding daily MMPA in June.

Discussion

Reasons for spatial differences

The increase of MMPA is in line with the mainstream measurement and model results (BARTHOLY, J. and PONGRÁTZ, R. 2007),




Fig. 7. Spearman correlation coefficients between elevation and different maximum precipitation amounts. M06 = June; M08 = August; p01 = 1%, p05 = 5% probability. Periods: 1960–1985 (red); 1986–2010 (blue)

although, PIRKHOFFER, E. et al. (2009) did not find an increase in the annual number of extreme precipitation events regarding Szeged and Budapest, except Debrecen in the 1901–2000 temporal range. PIRKHOFFER, E. et al. (2009) performed an analysis based on the annual dataset, thus, their findings can be different from our outcomes relating June and August. Our results are calculated from monthly data and this can be the explanation for the differences, too. They found that the majority of years had lower frequency of extreme events than the average. These longer periods were interrupted by short periods with a high number of extreme events.

In the countywide context, daily precipitation maximums in June showed hotspots. As the main wind direction is NW to SE the maximums are concentrated in the NW foothills of the Börzsöny Mountains and of the Western Carpathians, even though additional maxima appear in central TH and in the vicinity of Szeged. Conversely, in August, this trend is obscured probably by the occurrence of convective precipitation events in the centre of GHP. This theory is supported by the results of KENDON, E. et al. (2014) who concluded the highlighted role of convective precipitations in model efficiency in the summer. This is also in line with the results of MEZŐSI, G. et al. (2013), who modelled the highest increase in precipitation extremities on the GHP. In contrast, the correlation between MMPA and topography was strong in August and weak in June for the 1960–1985 period for the whole country.

Turning to the period 1986–2010 moderate to weak correlations were found without any structure in months or period length. This could partly be the result of spatial resolution, county borders are not identical with topographic elements. The general tendency is that most trends disappear due to the increasing extremities (GELYBÓ, Gy. et al. 2018). As a consequence of this, certain phenomena e.g. the occurrence of convective precipitation events would be less typical for one particular time period or location and diversity is expected to increase.

Risky and neutral areas of changes in intense rainfalls

Most differences in MMPA changes between the macro regions were significant in June and August pointing to regional variations of heavy rainfalls. Regarding the 6-days long MMPA in June the lowest increase was detected on the GHP. The maximum increase was in Nyírség. Since Nyírség already had the highest values (100 mm <) the most extreme 6-days precipitation conditions are presumed to appear here. However, in the Nyírség, the ratio of forest land cover is high (NÉGYESI, G. et al. 2015; NÉGYESI, G. 2018) and the soil texture is sandy (PÁSZTOR, L. et al. 2016a,b), the probability of pluvial flood occurrence remains still low. As in this region the most dangerous soil degrading factor is wind erosion, an increasing soil moisture content in early summer could provide defense against deflation (NÉGYESI, G. et al. 2016; PÁSZTOR, L. et al. 2016c). In contrast, at the vicinity of Szeged the highest increase, parallel with the highest precipitation volumes, triggers increasing pluvial flood risk as a result of loamy soil texture and of the highly degraded structure of the soils due to intensive agricultural cultivation (VAN LEEUWEN, B. et al. 2013). The largest change in the northern part of GHP (approx. + 30 mm) increased the original amount of ~ 65 mm leading only to moderate hazard.

Concerning the western part of the country and especially the southern part of the Transdanubian Hills, these long-term heavy precipitation, particularly after a very wet winter and spring, can trigger mass movements. For instance, landslides, bank failures along the shore of Danube are presumed to be the results of the precipitation pattern and the role of water level fluctuations and geological structure is less important (ÚJVÁRI, G. et al. 2009). Increasing precipitation extremity is a potential hazard to river bank stability in southern Hungary. Some parts of the southern TH have the highest increase both in June and August whereas the annual precipitation amount does not change. This underlines the importance of

water storage in the soil. EEKHOUT, J.P.C. et al. (2018) presumed a considerable decline in soil water storage capacity due to climate change in Europe. Hence, surface pools are going to have a special role in this region.

In August, daily MMPA increases in southern part of the Transdanubian Hills, central Hungary and northern Hungary. Generally, these parts of the country had moderate MMPAs, except the already discussed southern part of the TH. In this region, the ratio of crop fields is around 30 per cent. In August, the crop fields after wheat harvest have no canopy cover protection against heavy showers and the surface is generally covered by soil crust that minimizes infiltration leading to high runoff volume. Due to the hilly landscape and the loose parent material (loess) this region (LÓKI, J. 2010) would be the hot spot of the country not just from hydrological aspect, but from the aspect of erosion as well. Moreover, the southern part of the TH was found to be extremely vulnerable to flash flood hazard (PIRKHOFFER, E. et al. 2009; LÓCZY, D. et al. 2012), which is in line with our results as flash floods are the direct consequences of intense rainfalls.

Conclusions

The Goda-method applied on the CC database was found an effective tool for MMPAs for the temporal comparison. The next step has to be the validation of the predictions using local scale measured or country scale modelled data. The approach ensured us to reveal temporal and spatial differences in the changes of the mean maximum precipitation amount. Effect size pointed to the largest differences, and only the changes in August had large effects considering the differences between 1961–1985 and 1986–2010 both for the 1 and 5 per cent probabilities.

Regarding the spatial pattern of the southern part of the Transdanubian Hills, this is presumed to be the most endangered location as a result of the changing hydraulic circumstances. In this region, the probability of flash flood,

soil erosion, and mass movements increases, whereas the environmental conditions such as loose parent material and the high percentage of crop fields also point to potential danger. Therefore, the protection strategy has to concentrate on this part of the country. However, more detailed spatial and temporal resolution is needed in order to gain more reliable data at local scale. More frequent flash flood phenomena and the increasing erosion require the establishment of a storm monitoring system with high temporal resolution measurements. This seems to be necessary to track the changing intensity rainfall events and to prepare an adequate conservation strategy.

Acknowledgements: The research was supported by the National Research, Development and Innovation Office (NKFIH 108755).

REFERENCES

- BALÁZS, B., BÍRÓ, T., DYKE, G.J., SINGH, S.K. and SZABÓ, Sz. 2018. Extracting water-related features using reflectance data and principal component analysis of Landsat images. *Hydrological Sciences Journal* 63. (2): 269–284. Doi: 10.1080/02626667.2018.1425802
- BALLABIO, C., BORELLI, P., SPINONI, J., MEUSBURGER, K., MICHAELIDES, S., BEGUERIA, S., KLIK, A., PETAN, S., JANEČEK, M., OLSEN, P., AALTO, J., LAKATOS, M., RYMSZEWICZ, A., DUMITRESCU, A., PERČEC TADIĆ, M., DIODATO, M., KOSTALOVA, J., ROUSSEVA, S., BANASIK, K., ALEWELL, C. and PANAGOS, P. 2017. Mapping monthly rainfall erosivity in Europe. *Science of the Total Environment* 579. 1298–1315.
- BARTHOLY, J. and PONGRÁCZ, R. 2007. Regional analysis of extreme temperature and precipitation indices for the Carpathian Basin from 1946 to 2001. *Global and Planetary Change* 57. (1–2): 83–95.
- CHEVAL, S., DUMITRESCU, A.M. and BIRSAN, V. 2016. Variability of the aridity in the South Eastern Europe over 1961–2050. *Catena* 151. 74–86
- CZIGÁNY, Sz., PIKHOFFER, E. and GERESDI, I. 2010. Impact of extreme rainfall and soil moisture on flash flood generation. *Időjárás* 114. (1–2): 79–100.
- DÓVÉNYI, Z. (ed.) 2010. *Magyarország kistájainak katasztere* (Inventory of microregions in Hungary). Budapest, MTA Földrajztudományi Kutatóintézet.
- ECKHOUT, J.P.C., HUNINK, J.E., TERINK, W., DE VENITE, J. and SAN DIEGO, C. 2018. Why increased extreme precipitation under climate change negatively affects water security. *Hydrology and Earth System Sciences* 22. 5935–5946.
- FIELD, A. 2009. *Discovering statistics*. London, SAGE Publications.
- FOWLER, H.J. and KILSBY, C.G. 2003. Implications of changes in seasonal and annual extreme rainfall. *Geophysical Research Letters* 30. (13): 17–20. Available at <https://doi.org/10.1029/2003GL017327>
- GELYBÓ, Gy., TÓTH, E., FARKAS, Cs., HOREL, Á., KÁSA, I. and BAKACSI, Zs. 2018. Potential impacts of climate change on soil properties. *Agrokémia és Talajtan* 67. (1): 121–141.
- GODA, L. 1966. *Frequency of several days long precipitations*. Tanulmányok és kutatási eredmények 20. Budapest, VITUKI, 6–15. (in Hungarian)
- HOTHORN, T., HORNIK, K., VAN DE WIEL, M.A. and ZEILEIS, A. 2008. Implementing a class of permutation tests: The coin package. *Journal of Statistical Software* 28. (8): 1–23. URL <http://www.jstatsoft.org/v28/i08/>
- IHÁSZ, I., MÁTRAI, A., SZINTAI, B., SZÚCS, M. and BONTA, I. 2018. Application of European numerical weather prediction models for hydrological purposes. *Időjárás* 122. (1): 59–79. Doi:10.28974/idojaras.2018.1.5
- JAKAB, G., MADARÁSZ, B., SZABÓ, J.A., TÓTH, A., ZACHÁRY, D., SZALAI, Z., KERTÉSZ, Á. and DYSON, J. 2017. Infiltration and soil loss changes during the growing season under ploughing and conservation tillage. *Sustainability* 9. (1726): 1–13. Doi: 10.3390/su9101726
- JAKAB, G., SZABÓ, J. and SZALAI, Z. 2015. A review on sheet erosion measurements in Hungary. *Journal of Landscape Ecology* 13. (1): 89–103.
- KENDON, E.J., ROBERTS, N.M., FOWLER, H.J., ROBERTS, M.J., CHAN, S.C. and SENIOR, C.A. 2014. Heavier summer downpours with climate change revealed by weather forecast resolution model. *Natural Climate Change* 4. (7): 570–576.
- KERTÉSZ, Á. and CENTERI, Cs. 2006. Hungary. In *Soil erosion in Europe*. Eds.: BOARDMAN, J. and POESEN, J., Chichester, John Wiley & Sons, 139–153.
- KNAPP, A.K., BEIER, C., BRISKE, D.D., CLASSEN, A.T., LUO, Y., REICHSTEIN, M., SMITH, M.D., SMITH, S.D., BELL, J.E., FAY, P.A., HEISLER, J.L., LEAWITT, S.V., SHERRY, R., SMITH, B. and WENG, E. 2008. Consequences of more extreme precipitation regimes for terrestrial ecosystems. *BioScience* 58. (9): 811–821. Available at <https://doi.org/10.1641/B580908>
- KRISTÓF, E., PONGRÁCZ, R. and BARTHOLY, J. 2017. Távkapcsolati rendszerek hatása a Kárpát-medence térségére (Impact of distant connection systems on the surroundings of the Carpathian Basin). In *HUNGEO 2017. Bányászat és környezet – harmóniában*. Eds.: CSERNY, T. and ALPEK, B.L., Budapest, Magyarhoni Földtani Társulat, 75–76.
- LAKATOS, M. and HOFFMANN, L. 2018. Növekvő csapadékinzultás, magasabb mértékadó csapadékok a változó klímában (Increasing precipitation intensity, higher standard precipitations in changing climate). In *Országos Települési Csapadékvíz-gazdálkodási Konferencia Tanulmányai*. Ed.: BÍRÓ, T., Budapest, Dialóg Campus Kiadó, 8–16.

- LÁSZLÓ, E. and SALAVEC, P. 2018. Relationship between weather conditions advantageous for the development of urban heat island and atmospheric macro-circulation changes. *International Journal of Climatology* 38. 3224–3232.
- LÁSZLÓ, E., BOTTYÁN, Z. and SZEGEDI, S. 2016. Long-term changes of meteorological conditions of urban heat island development in the region of Debrecen, Hungary. *Theoretical and Applied Climatology* 124. (1): 365–373.
- LÓCZY, D., CZIGÁNY, SZ. and PIRKHOFFER, E. 2012. Flash flood hazards. In *Studies on Water Management Issues*. Ed.: KUMARASAMY, M., Rijeka, InTech, 57–72. Doi: 10.5772/28775
- LÓKI, J. 2010. Agriculture: Crop Cultivation and Horticulture. In *Anthropogenic Geomorphology: A Guide to Man-Made Landforms*. Eds.: SZABÓ, J., DÁVID, L. and LÓCZY, D., Dordrecht–Heidelberg–New York, Springer, 55–71.
- LOVE, J. and MAIR, P. 2017. *Walrus: Robust Statistical Methods*. (R package version 1.0.1.). Available at <https://cran.r-project.org/package=walrus>
- MAHERAS, P., TOLIKÁ, K., ANAGNOSTOPOULOU, C., MAKRA, L., SZPIROSZ, K. and KÁROSSY, CS. 2018. Relationship between mean and extreme precipitation and circulation types over Hungary. *International Journal of Climatology* 38. (12): 4518–4532. Doi: 10.1002/joc.5684
- MEZŐSI, G., MEYER, B.C., LOIBL, W., AUBRECHT, C., CSORBA, P. and BATA, T. 2013. Assessment of regional climate change impacts on Hungarian landscapes. *Regional Environmental Change* 13. (4): 797–811. Doi:10.1007/s10113-012-0326-1
- MÍKA, J. and FARKAS, A. 2017. Sensitivity of inland water bodies, natural vegetation and agriculture to weather extremes and to climate change. *Tájökológiai Lapok* 15. (2): 85–90. (in Hungarian with English abstract)
- MÍLOSEVIC, D. and SAVIC, S. 2013. Analysis of precipitation quantities and trends from Pannonian and Peripannonian parts of Serbia. *Dela* 39. 125–139.
- MÍLOSEVIC, D., SAVIC, S., PANTELIC, M., STANKOV, U., ZIBERNA, I., DOLINAJ, D. and LESCESEN, I. 2017. Variability of seasonal and annual precipitation in Slovenia and its correlation with large-scale atmospheric circulation. *Open Geosciences* 8. (1): 593–605.
- MÜLLER, M., KAŠPAR, M. and MATSCHULLAT, J. 2009. Heavy rains and extreme rainfall-runoff events in Central Europe from 1951 to 2002. *Natural Hazards and Earth System Sciences* 9. 441–450. Available at <https://doi.org/10.5194/nhess-9-441-2009>
- NÉGYESI, G. 2018. Surveying the spatiotemporal changes of afforestation in the Nyírség – from the aspect of wind erosion. *Tájökológiai Lapok* 16. (2): 113–128. (in Hungarian with English abstract)
- NÉGYESI, G., LÓKI, J., BURÓ, B. and SZABÓ, SZ. 2016. Effect of soil parameters on the threshold wind velocity and maximum eroded mass in a dry environment. *Arabian Journal of Geosciences* 9. 588–599.
- NÉGYESI, G., LÓKI, J., BURÓ, B., SZABÓ, J., BAKACSI, ZS. and PÁSZTOR, L. 2015. The potential wind erosion map of an area covered by sandy and loamy soils – based on wind tunnel measurements. *Zeitschrift für Geomorphologie* 59. (1): 59–77.
- PANAGOS, P., BALLABIO, C., BORELLI, P., MEUSBURGER, K., KLIK, A., ROUSSEVA, S., PERČEC TADIĆ, M., MICHAELIDES, S., HRABÁLKOVÁ, M., OLSEN, P., AALTO, J., LAKATOS, M., RYMSZEWICZ, A., DUMITRESCU, A., BEGUERIA, S. and ALEWELL, C. 2015. Rainfall erosivity in Europe. *Science of the Total Environment* 511. 801–814.
- PÁSZTOR, L., KÖRÖSPARTI, J., BOZÁN, CS., LABORCZI, A. and TAKÁCS, K. 2015. Spatial risk assessment of hydrological extremities: Inland excess water hazard, Szabolcs-Szatmár-Bereg county, Hungary. *Journal of Maps* 11. (4): 636–644.
- PÁSZTOR, L., NÉGYESI, G., LABORCZI, A., KOVÁCS, T., LÁSZLÓ, E. and BIHARI, Z. 2016a. Integrated spatial assessment of wind erosion risk in Hungary. *Natural Hazards and Earth System Sciences* 16. 2421–2432.
- PÁSZTOR, L., LABORCZI, A., TAKÁCS, K., SZATMÁRI, G., ILLÉS, G., FODOR, N., NÉGYESI, G., BAKACSI, ZS. and SZABÓ, J. 2016b. Spatial distribution of selected soil features in Hajdú-Bihar county represented by digital soil maps. *Acta Geographica Debrecina. Landscape and Environment* 10. (3–4): 203–213.
- PÁSZTOR, L., WALTNER, I., CENTERI, CS., BELÉNYESI, M. and TAKÁCS, K. 2016c. Soil erosion of Hungary assessed by spatially explicit modelling. *Journal of Maps* 12. (1): 407–414. Doi: 10.1080/17445647.2016.1233913
- PIECZKA, I., PONGRÁCZ, R. and BARTHOLY, J. 2011. Expected trends of regional climate change for the Carpathian Basin for the 21st century. *International Journal of Environment and Pollution* 46. (1–2): 6–17.
- PIRKHOFFER, E., CZIGÁNY, SZ. and GERESDI, I. 2009. Impact of rainfall pattern on the occurrence of flash floods in Hungary. *Zeitschrift für Geomorphologie N. F.* 53. Supplement 2. 139–157.
- PONGRÁCZ, R., BARTHOLY, J. and KIS, A. 2014. Estimation of future precipitation conditions for Hungary with special focus on dry periods. *Időjárás* 118. (4): 305–321.
- R. Core Team 2017. *R: A Language and Environment for Statistical Computing*. Vienna, R Foundation for Statistical Computing. Available at <https://www.R-project.org/>
- RENARD, K.G. and FREIMUND, J.R. 1994. Using monthly precipitation data to estimate the R-factor in the revised USLE. *Journal of Hydrology* 157. (1–4): 287–306.
- RODRIGO-COMINO, J., NEUMANN, M., REMKE, A. and RIES, J.B. 2019. Assessing environmental changes in abandoned German vineyards. Understanding key issues for restoration management plans. *Hungarian Geographical Bulletin* 67. (4): 319–332.
- SPINONI, J., LAKATOS, M., SZENTIMREY, T., BIHARI, Z., SZALAI, S., VOGT, J. and ANTOFIE, T. 2015. Heat and cold waves trends in the Carpathian Region from

- 1961 to 2010. *International Journal of Climatology* 35. (14): 4197–4209. Doi:10.1002/joc.4279
- SULLIVAN, G.M. and FEINN, R. 2012. Using effect size – or why the P value is not enough. *Journal of Graduate Medical Education* 4. (3): 279–282. Doi: 10.4300/JGME-D-12-00156.1
- SZALAI, S., AUER, I., HIEBL, J., MILKOVICH, J., RADIM, T., STEPANEK, P., ZAHRADNICEK, P., BIHARI, Z., LAKATOS, M., SZENTIMREY, T., LIMANOWKA, D., KILAR, P., CHEVAL, S., DEÁK, GY., MIHIC, D., ANTOLOVIC, I., NEJEDLIK, P., STASTNY, P., MIKULOVA, K., NABYVANETS, I., SKYRYK, O. and KRAKOVSKAYA, S. 2013. *Climate of the Greater Carpathian Region. Final technical report*. Available at <http://www.carpatclim-eu.org>
- SZENTIMREY, T. 2011. *Manual of homogenization software MASHv3. 03*. Budapest, Hungarian Meteorological Service.
- SZENTIMREY, T. and BIHARI, Z. 2007. Mathematical background of spatial interpolation, Meteorological interpolation based on surface homogenized data bases (MISH). In *COST Action 719*. Budapest, COST Office, 17–27.
- SZENTIMREY, T., LAKATOS, M., BIHARI, Z., KOVACS, T., SZALAI, S., AUER, I., HIEBL, J., MILKOVIC, J., STEPANEK, P. and ZAHRADNICEK, P. 2012. *Final Report on Quality Control and Data Homogenization Measures Applied per Country, Including QC Protocols and Measures to Determine the Achieved Increase in Data Quality*. 12. CARPATCLIM Project Deliverable D1. Available at http://www.CARPATCLIM-eu.org/docs/deliverables/D1_12.pdf
- SZÚCS, P., CSEPINSZKY, B., SISÁK, I. and JAKAB, G. 2006. Rainfall simulation in wheat culture at harvest. *Cereal Research Communications* 34. (1): 81–84.
- ÚJVÁRI, G., MENTES, GY., BÁNYAI, L., KRAFT, J., GYIMÓTHY, A. and KOVÁCS, J. 2009. Evolution of a bank failure along the River Danube at Dunaszekcső, Hungary. *Geomorphology* 109. 197–209.
- VAN LEEUWEN, B., HENITS, L., MÉSZÁROS, M., SZATMÁRI, J., TOBAK, Z., PAVIĆ, D., SAVIĆ, S. and DOLINAJ, D. 2013. RapidEye satellite imagery for inland excess water identification. *Hidrológiai Közlöny* 93. (3): 17–24. (in Hungarian with English abstract)
- WALTNER, I., PÁSZTOR, L., CENTERI, Cs., TAKÁCS, K., PIRKÓ, B., KOÓS, S. and LÁSZLÓ, P. 2018. Evaluating the new soil erosion map of Hungary – a semi-quantitative approach. *Land Degradation & Development* 29. (4): 1295–1302.
- WICKHAM, H. and HENRY, L. 2018. *Tidyr: Easily tidy data with 'spread ()' and 'gather ()' functions*. R package version 0.8.0. Available at <https://CRAN.R-project.org/package=tidyr>
- WISCHMEIER, W.H. and SMITH, D.D. 1978. *Predicting rainfall erosion losses: A guide to conservation planning*. USDA Agricultural Handbook 537. Washington D. C., US Government Printing Office.
- ZHANG, F., ODINS, A.M. and NIELSEN-GAMMON, J.W. 2006. Mesoscale predictability of an extreme warm-season precipitation event. *Weather and Forecasting* 21. 149–166. Available at <https://doi.org/10.1175/WAF909.1>

